

ALEMTUZUMAB (SUBCUTANEOUS)
(CAMPATH™)

REVISED January 2014

Date written _____ To begin _____

Patient's: Height _____ cm Weight _____ kg BSA _____ m²
Allergies: No Yes: _____ Diagnosis _____ Metastatic Site _____

1. Check that:

- CBC, Diff & Plts, Bun/Creat, Uric Acid have been done recently
- Patient is taking allopurinol
- Weekly CBC with Diff & Plts, Bun/Creat. If ANC <250 and/or Plts <25,000, hold Rx, call MD.
If ANC or Plts decrease to <50% baseline level hold Rx, call MD
- Serum Electrolytes & LFT's q 3 weeks

2. Consent Obtained?

- Yes
- Preprinted Consent

3. Infusion Room General Order Set will be initiated

4. Premedicate 30 minutes prior to injection:

Diphenhydramine 25 mg po
Acetaminophen 1,000 mg po

Continue premeds until patient has received 4 doses Alemtuzumab 30 mg, then D/C

5. Chemotherapy:

Day 1 Alemtuzumab 3 mg sub-Q
Day 3 Alemtuzumab 10 mg sub-Q
Day 5 Alemtuzumab 30 mg sub-Q (dose divided into 2 injections)

**Continue Alemtuzumab 30 mg sub-Q qMon, Wed, Fri to continue until
day _____, date: _____. Initials _____ MD.**

**If therapy is interrupted for > 7 days due to hematologic toxicity
(severe neutropenia or thrombocytopenia) reinitiate therapy at:**

3 mg sub-Q (if first occurrence of toxicity) and dose escalate as above
10 mg sub-Q (if second occurrence of toxicity) and dose escalate as above

If third occurrence of toxicity, contact MD & consider discontinuation of therapy

Premedications do not need to be given if patient has been in treatment for > 2 weeks

6. Prophylactic antibiotics at discharge

Trimethoprim/Sulfamethoxazole prophylaxis (requires Prescription)
Acyclovir (requires prescription)

Preparer's Signature _____ Date _____

Attending's Signature _____ Printed name _____ Pager number _____ Date _____ Time AM/PM _____



MR CHEMO ORDER